

Cluster Production in pBUU - Past and Future

Pawel Danielewicz

National Superconducting Cyclotron Laboratory
Michigan State University

Transport 2017: International Workshop
on Transport Simulations for Heavy Ion Collisions
under Controlled Conditions

FRIB-MSU, East Lansing, Michigan, March 27 - 30, 2017



Boltzmann Equation Model (BEM/pBUU)

Degrees of freedom (X):

nucleons, deuterons, tritons, helions ($A \leq 3$), Δ , N^* , pions

Fundamentals:

- Relativistic Landau theory (Chin/Baym)
Energy functional (ϵ)
- Real-time Green's function theory
Production/absorption rates ($\mathcal{K}^<$, $\mathcal{K}^>$)

$$\frac{\partial f}{\partial t} + \frac{\partial \epsilon}{\partial \mathbf{p}} \frac{\partial f}{\partial \mathbf{r}} - \frac{\partial \epsilon}{\partial \mathbf{r}} \frac{\partial f}{\partial \mathbf{p}} = \mathcal{K}^< (1 \mp f) - \mathcal{K}^> f$$

production

absorption rate



Single-Particle Energies & Functional

$$\frac{\partial f}{\partial t} + \frac{\partial \epsilon}{\partial \mathbf{p}} \frac{\partial f}{\partial \mathbf{r}} - \frac{\partial \epsilon}{\partial \mathbf{r}} \frac{\partial f}{\partial \mathbf{p}} = \mathcal{K}^< (1 \mp f) - \mathcal{K}^> f$$

The single-particle energies ϵ are given in terms of the net energy functional $E\{f\}$ by,

$$\epsilon(\mathbf{p}) = \frac{\delta E}{\delta f(\mathbf{p})}$$

In the local cm, the mean potential is

$$U_{opt} = \epsilon - \epsilon_{kin}$$

and $\epsilon_{kin} = \sqrt{p^2 + m^2}$



Energy Functional

The functional:

$$E = E_{vol} + E_{gr} + E_{iso} + E_{Coul}$$

where

$$E_{gr} = \frac{a_{gr}}{\rho_0} \int d\mathbf{r} (\nabla \rho)^2$$

For covariant volume term, ptcle velocities parameterized in local frame:

$$v^*(p, \rho) = \frac{p}{\sqrt{p^2 + m^2} \left(1 + c \frac{\rho}{\rho_0} \frac{1}{(1 + \lambda p^2/m^2)^2} \right)^2}$$

precluding a supraluminal behavior, with ρ - baryon density. The 1-ptcle energies are then

$$\epsilon(p, \rho) = m + \int_0^p dp' v^* + \Delta\epsilon(\rho)$$


Parameters in the velocity varied to yield different optical potentials characterized by values of effective mass,

$$m^* = p_F / v_F.$$



Structure Interface

Potential from
p-scattering
(Hama *et al.*
PRC41(90)2737)
&
parameterizations




Ground-state densities from
electron scattering and from
functional minimization.

From $E(f) = \min$:

$$0 = \epsilon \left(p^F(\rho) \right) - 2 a_{gr} \nabla^2 \left(\frac{\rho}{\rho_0} \right) - \mu$$

⇒ **Thomas-Fermi eq.**



Many-Body Theory

Transport eq. for nucleons follows from the eq. of motion for the 1-ptcle Green's function (KB eq.). Transport eq. for deuterons ($A = 2$) from the eq. for 2-ptcle Green's function??

Wigner function in second quantization

$$f(\mathbf{p}; \mathbf{R}, T) = \int d\mathbf{r} e^{-i\mathbf{p}\mathbf{r}} \langle \hat{\psi}_H^\dagger(\mathbf{R} - \mathbf{r}/2, T) \hat{\psi}_H(\mathbf{R} + \mathbf{r}/2, T) \rangle$$

where $\langle \cdot \rangle \equiv \langle \Psi | \cdot | \Psi \rangle$ and $|\Psi\rangle$ describes the initial state.


Evolution driven by a Hamiltonian. Interaction Hamiltonian:

$$\hat{H}^1 = \frac{1}{2} \int d\mathbf{x} d\mathbf{y} \hat{\psi}^\dagger(\mathbf{x}) \hat{\psi}^\dagger(\mathbf{y}) v(\mathbf{x} - \mathbf{y}) \hat{\psi}(\mathbf{y}) \hat{\psi}(\mathbf{x}),$$



Evolution Contour

$$\begin{aligned}
 \langle \hat{O}_H(t_1) \rangle &= \langle T^a \left[\exp \left(-i \int_{t_1}^{t_0} dt' \hat{H}_I^1(t') \right) \right] \hat{O}_I(t_1) \\
 &\quad \times T^c \left[\exp \left(-i \int_{t_0}^{t_1} dt' \hat{H}_I^1(t') \right) \right] \rangle \\
 &= \langle T \left[\exp \left(-i \int_{t_0}^{t_0} dt' \hat{H}_I^1(t') \right) \hat{O}_I(t_1) \right] \rangle,
 \end{aligned}$$



Expectation value expanded perturbatively in terms of V and noninteracting 1-ptcle Green's functions on the contour

$$iG_0(\mathbf{x}, t, \mathbf{x}', t') = \langle T \left[\hat{\psi}_I(\mathbf{x}, t) \hat{\psi}_I^\dagger(\mathbf{x}', t') \right] \rangle$$




Single-Particle Evolution

Wigner function corresponds to a particular case of the Green's function on contour:


$$f(\mathbf{p}; \mathbf{R}, T) = \int d\mathbf{r} e^{-i\mathbf{p}\mathbf{r}} (\mp i) G^<(\mathbf{R} + \mathbf{r}/2, T, \mathbf{R} - \mathbf{r}/2, T)$$

If we find an equation for G , this will also be an equation for f .

Dyson eq. from perturbation expansion:



$$G = G_0 + G_0 \Sigma G$$



Outcome of Evolution

Formal solution of the Dyson eq:

$$\mp iG^<(x, t; x', t') = \int dx_1 dt_1 dx'_1 dt'_1 G^+(x, t; x_1, t_1) \\ \times (\mp i)\Sigma^<(x_1, t_1; x'_1, t'_1) G^-(x, t; x_1, t_1)$$

and

$$\mp i\Sigma^<(x, t; x', t') = \langle \hat{j}^\dagger(x', t') \hat{j}(x, t) \rangle_{\text{irred}}$$

where the source j is

$$\hat{j}(x, t) = [\hat{\psi}(\mathbf{x}, t), \hat{H}^1]$$



Quasiparticle Limit

Under slow spatial and temporal changes in the system, the Green's function expressible in terms of the Wigner function f and 1-ptcle energy ϵ_p

$$\mp iG^<(x, t; x', t') \approx \int dp f(p; \frac{x+x'}{2}, \frac{t+t'}{2}) e^{i(p(x-x') - \epsilon_p(t-t'))}$$

Then also Boltzmann eq:

$$\frac{\partial f}{\partial t} + \frac{\partial \epsilon_p}{\partial \mathbf{p}} \frac{\partial f}{\partial \mathbf{r}} - \frac{\partial \epsilon_p}{\partial \mathbf{r}} \frac{\partial f}{\partial \mathbf{p}} = -i\Sigma^<(1-f) - i\Sigma^>f$$

$$\mp i\Sigma^< : \quad \begin{array}{c} | \\ \bullet \\ \swarrow \quad \searrow \\ \nearrow \quad \nwarrow \\ \bullet \\ | \end{array} = \left| \begin{array}{c} \swarrow \quad \nearrow \\ \bullet \\ \nwarrow \quad \searrow \end{array} \right|^2 \times \begin{array}{c} | \\ \rightarrow \\ \leftarrow \\ \rightarrow \\ | \end{array}$$

2-Particle Green's Function

Transport eq. for deuterons ($A = 2$) from the eq. for 2-ptcle Green's function??

$$iG_2^< = \langle \hat{\psi}^\dagger(\mathbf{x}'_1 t') \hat{\psi}^\dagger(\mathbf{x}'_2 t') \hat{\psi}(\mathbf{x}_2 t) \hat{\psi}(\mathbf{x}_1 t) \rangle$$

For the contour function:

$$G_2 = \mathcal{G} + \mathcal{G} v G_2$$

where \mathcal{G} – irreducible part of G_2 (w/o two 1-ptcle lines connected by the potential v ; anything else OK)

In terms of retarded Green's function $G_2^<$:

$$iG_2^< = (1 + v G_2^+) i\mathcal{G}^< (1 + v G_2^-)$$



Deuteron Quasiparticle Limit

In the limit of slow spatial and temporal changes, deuteron contribution to the 2-ptcle Green's function:

$$\begin{aligned}
 iG_2^< &= \langle \hat{\psi}^\dagger(\mathbf{x}'_1 t') \hat{\psi}^\dagger(\mathbf{x}'_2 t') \hat{\psi}(\mathbf{x}_2 t) \hat{\psi}(\mathbf{x}_1 t) \rangle \\
 &\simeq \int d\mathbf{p} f_d(\mathbf{p} \mathbf{R} T) \phi_d^*(r') \phi_d(r) e^{i\mathbf{p} \cdot \left(\frac{\mathbf{x}_1 + \mathbf{x}_2}{2} - \frac{\mathbf{x}'_1 + \mathbf{x}'_2}{2} \right)} e^{-i\epsilon_d(t-t')} \\
 &\quad + \dots,
 \end{aligned}$$

where $\mathbf{R} = \frac{1}{4}(\mathbf{x}_1 + \mathbf{x}_2 + \mathbf{x}'_1 + \mathbf{x}'_2)$, $\mathbf{r} = \mathbf{x}_1 - \mathbf{x}_2$

ϕ_d and f_d – internal wave function and cm Wigner function

$\dots \equiv$ continuum

Transport eq from integral quantum eq of motion:

$$\frac{\partial f_d}{\partial T} + \frac{\partial \epsilon_d}{\partial \mathbf{p}} \frac{\partial f_d}{\partial \mathbf{R}} - \frac{\partial \epsilon_d}{\partial \mathbf{R}} \frac{\partial f_d}{\partial \mathbf{p}} = \mathcal{K}^< (1 + f_d) - \mathcal{K}^> f_d$$




Wave Equation

From Green's function eq, the equation for wavefunction:

$$(\epsilon_d(\mathbf{P}) - \epsilon_N(\mathbf{P}/2 + \mathbf{p}) - \epsilon_N(\mathbf{P}/2 - \mathbf{p})) \phi_d(\mathbf{p}) - (1 - f_N(\mathbf{P}/2 + \mathbf{p}) - f_N(\mathbf{P}/2 - \mathbf{p})) \int d\mathbf{p}' v(\mathbf{p} - \mathbf{p}') \phi_d(\mathbf{p}') = 0$$

In zero-temperature matter, discrete states lacking over a vast range of momenta



Cluster Production & Absorption

?? Production & absorption rates: $i\mathcal{K}^> = \phi^* v i\mathcal{G}^> v \phi$

Leading contribution

$$\mathcal{K}^< = \int d\mathbf{r} d\mathbf{r}' \phi_d^* v \langle \hat{\psi}^\dagger(\mathbf{x}'_1 t') \hat{\psi}(\mathbf{x}_1 t) \rangle \langle \hat{\psi}^\dagger(\mathbf{x}'_2 t') \hat{\psi}(\mathbf{x}_2 t) \rangle v \phi_d$$

Leading-order in the quasiparticle expansion: neutron & proton come together and make a deuteron.


If system approximately uniform and stationary, the process not allowed by energy-momentum conservation.

Process possible in a mean field varying in space, but, in nuclear case, the high-energy production rate low – tested in Glauber model.




3-Nucleon Collisions

production
amplitude



d made in a 3N collision



First correction to the pure 1-ptcle state, from a coupling to p-h excitations, yields a contribution to the d-production due to 3-nucleon collisions.

Still more nucleons involved in production of heavier clusters.



Deuteron Production

Detailed balance:

$$|\mathcal{M}^{npN \rightarrow Nd}|^2 = |\mathcal{M}^{Nd \rightarrow Nnp}|^2 \propto d\sigma^{Nd \rightarrow Nnp}$$



Thus, production can be described in terms of breakup.


Problem: Breakup cross section only known over limited range of final states - Interpolation/extrapolation needed

Impulse approximation works at high incident energy

$$|\overline{\mathcal{M}}_{Nd \rightarrow Npn}|_{IA}^2 = \left| \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \end{array} \right|^2 + \left| \begin{array}{c} \text{Diagram 4} \\ \text{Diagram 5} \\ \text{Diagram 6} \end{array} \right|^2 + \left| \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \\ \text{Diagram 9} \end{array} \right|^2$$

Renormalized Impulse Approximation


Renormalization factor for squared matrix element to get breakup cross section right as a function of energy



$$d\sigma^{Nd \rightarrow Nnp} \propto F \sigma_{NN} |\phi_d(p)|^2$$




Single-Particle Spectra



proton

&



deuteron inclusive spectra

histograms: calculations using

$$|\mathcal{M}^{npN \rightarrow Nd}|^2 = |\mathcal{M}^{Nd \rightarrow npN}|^2 \propto d\sigma_{Nd \rightarrow npN}$$

and $\langle f \rangle < 0.2$ cut-off for deuterons



A = 3 Particles + Tests


A=3-ptcles from 4N collisions

Christiane Kuhrts: solving finite- T Galitski-Feynman (GF) and modified (in-medium) Alt-Grassberger-Sandhas eqs

solid lines: finite- T GF for cross-sections and existence

dashed lines: free cross sections + $\langle f \rangle$ cut-off

symbols: INDRA data $^{129}\text{Xe} + ^{119}\text{Sn}$ at 50 MeV/nucleon



Cluster Yields and Entropy

Compression in central reactions accompanied by heating. Is the matter heated as much as expected for shock compression??

Experimental measure of entropy: relative cluster yields

$$E = TS - PV + \mu A \quad \Leftrightarrow \quad 3AT/2 \simeq TS - AT + \mu A$$

as at freeze-out ideal gas and then


$$\frac{S}{A} \simeq \frac{5}{2} - \frac{\mu}{T}$$

In equilibrium

$$\frac{N_d}{N_p} \propto \frac{\exp\left(\frac{2\mu}{T}\right)}{\exp\left(\frac{\mu}{T}\right)} \quad \Rightarrow \quad \frac{S}{A} \simeq 3.9 - \log\left(\frac{N_d}{N_p}\right)$$



Validity of Entropy Determination



Collective Expansion

Au+Au 250A MeV $60^\circ < \theta_{cm} < 90^\circ$


Is expansion viscous or isentropic?? Is pressure carrying out work producing a collective expansion of matter?

$$\langle E_x \rangle = \frac{3}{2} T + \frac{m_x \langle v^2 \rangle}{2}$$


$$= \frac{3}{2} T + A_x \frac{m_N \langle v^2 \rangle}{2}$$

In isentropic expansion, average kinetic energy should increase with fragment mass.

yield
ratio



av kin
energy



EXP(erat5) Tr.Mod.(b<3.5fm)

■ p	□ p
◆ d	◇ d
● t	○ t
▲ ³ He	△ ³ He
▼ ⁴ He	


filled symbols
-- data, Poggi et al
open symbols
-- calculation

Energy increases linearly!



Head-On Au + Au (FOPI)

Rapidity Distribution





400 MeV/nucleon



$A = 3$ in Head-On Au + Au (FOPI)


Rapidity Distributions



400 MeV/nucleon





Semicentral Au + Au (FOPI)



Semicentral Au + Au (FOPI)

Elliptic Flow



400 MeV/nucleon




Future of Light-Cluster Production in Transport

Production rate for cluster of mass A :

$$\begin{aligned} \mathcal{K}^<(\mathbf{p}_A) = & \int d\mathbf{p}'_1 \dots d\mathbf{p}'_{N'} d\mathbf{p}_1 \dots d\mathbf{p}_{N-1} |\mathcal{M}_{1'+\dots+N' \rightarrow 1+\dots+A}|^2 \\ & \times \delta(\mathbf{p}'_1 + \dots + \mathbf{p}'_{N'} - \mathbf{p}_1 - \dots - \mathbf{p}_{N-1} - \mathbf{p}_A) \\ & \times \delta(\epsilon'_1 + \dots + \epsilon'_{N'} - \epsilon_1 - \dots - \epsilon_{N-1} - \epsilon_A) \\ & \times f_1 \dots f_{N'} (1 \pm f_1) \dots (1 \pm f_{N-1}) \end{aligned}$$

Determination and sampling of separate $|\mathcal{M}|^2$ for every possible process. . . Potential nightmare! E.g.




Any simplifications??



Simplified Matrix Elements

Batko, Randrup, Vetter
NPA536(92)786

$|\mathcal{M}|^2 \propto 1 \Rightarrow$ Mini Fireball
??Too much dissipation??



Mini
Compound
Nucleus

Generalized coalescence:

$$|\mathcal{M}|^2 \propto \theta(p_0 - |\frac{p_A}{A} - p'_1|) \cdots \theta(p_0 - |\frac{p_A}{A} - p'_{N'}|)$$

Branching??

Automation needed!



Conclusions

- Real-time many-body theory provides fundamentals for production of clusters in transport theory
- Few-body collisions or rapidly changing mean-field conditions are needed to spur cluster production
- Detailed balance must be obeyed for thermodynamic consistency
- Breakup data yield production rates in collisions
- Clusters emphasize collective motion and provide information on phase-space densities and entropy
- Production description needs to be simplified in extending reach of theory.

Supported by National Science Foundation under Grant US PHY-1403906



Conclusions

- Real-time many-body theory provides fundamentals for production of clusters in transport theory
- Few-body collisions or rapidly changing mean-field conditions are needed to spur cluster production
- Detailed balance must be obeyed for thermodynamic consistency
- Breakup data yield production rates in collisions
- Clusters emphasize collective motion and provide information on phase-space densities and entropy
- Production description needs to be simplified in extending reach of theory.

Supported by National Science Foundation under Grant US PHY-1403906



Conclusions

- Real-time many-body theory provides fundamentals for production of clusters in transport theory
- Few-body collisions or rapidly changing mean-field conditions are needed to spur cluster production
- Detailed balance must be obeyed for thermodynamic consistency
- Breakup data yield production rates in collisions
- Clusters emphasize collective motion and provide information on phase-space densities and entropy
- Production description needs to be simplified in extending reach of theory.

Supported by National Science Foundation under Grant US PHY-1403906



Conclusions

- Real-time many-body theory provides fundamentals for production of clusters in transport theory
- Few-body collisions or rapidly changing mean-field conditions are needed to spur cluster production
- Detailed balance must be obeyed for thermodynamic consistency
- Breakup data yield production rates in collisions
- Clusters emphasize collective motion and provide information on phase-space densities and entropy
- Production description needs to be simplified in extending reach of theory.

Supported by National Science Foundation under Grant US PHY-1403906



Conclusions

- Real-time many-body theory provides fundamentals for production of clusters in transport theory
- Few-body collisions or rapidly changing mean-field conditions are needed to spur cluster production
- Detailed balance must be obeyed for thermodynamic consistency
- Breakup data yield production rates in collisions
- Clusters emphasize collective motion and provide information on phase-space densities and entropy
- Production description needs to be simplified in extending reach of theory.

Supported by National Science Foundation under Grant US PHY-1403906



Conclusions

- Real-time many-body theory provides fundamentals for production of clusters in transport theory
- Few-body collisions or rapidly changing mean-field conditions are needed to spur cluster production
- Detailed balance must be obeyed for thermodynamic consistency
- Breakup data yield production rates in collisions
- Clusters emphasize collective motion and provide information on phase-space densities and entropy
- Production description needs to be simplified in extending reach of theory.

Supported by National Science Foundation under Grant US PHY-1403906



Conclusions

- Real-time many-body theory provides fundamentals for production of clusters in transport theory
- Few-body collisions or rapidly changing mean-field conditions are needed to spur cluster production
- Detailed balance must be obeyed for thermodynamic consistency
- Breakup data yield production rates in collisions
- Clusters emphasize collective motion and provide information on phase-space densities and entropy
- Production description needs to be simplified in extending reach of theory.

Supported by National Science Foundation under Grant US PHY-1403906

