





# Collective Vibrations



## Giant Dipole Resonance




**Figure 6-18** Total photoabsorption cross section for  $^{197}\text{Au}$ . The experimental data are from S. C. Fultz, R. L. Bramblett, J. T. Caldwell, and N. A. Kerr, *Phys. Rev.* **127**, 1273 (1962). The solid curve is of Breit-Wigner shape with the indicated parameters.




# Particle-Hole excitations

Hartree-Fock state



$$|HF\rangle$$


1 particle-1 hole (1p1h) state




$$a_p^\dagger a_h |HF\rangle$$

2 particle-2 hole (2p2h) state

$$a_p^\dagger a_{p'}^\dagger a_h a_{h'} |HF\rangle$$





スライド: 松柳研一氏

# Tamm-Dancoff Approximation

$$\begin{aligned}\text{Assume: } |\nu\rangle = Q_\nu^\dagger |HF\rangle &= \sum_{ph} X_{ph} a_p^\dagger a_h |HF\rangle \\ &= \sum_{ph} X_{ph} |ph^{-1}\rangle\end{aligned}$$

(superposition of 1p1h states)

$$H|\nu\rangle = E_\nu|\nu\rangle$$




$$\sum_{p'h'} H_{ph,p'h'} X_{p'h'} = E_\nu X_{ph}$$

residual  
interaction

$$H_{ph,p'h'} = (\epsilon_p - \epsilon_h) \delta_{ph,p'h'} + \langle ph' | \bar{v} | hp' \rangle$$

Tamm-Dancoff equation



スライド: 松柳研一氏

$$V(\mathbf{r}) \sim \int d\mathbf{r}' v(\mathbf{r}, \mathbf{r}') \rho(\mathbf{r}')$$

vibration:  $\rho = \rho_0(\mathbf{r}) \rightarrow \rho_0(\mathbf{r}) + \delta\rho(\mathbf{r}, t)$

---


↓  
residual  
interaction


# TDA on a schematic model


Separable interaction:  $\langle ph' | \bar{v} | hp' \rangle = \lambda D_{ph} D_{p'h'}^*$

Tamm-Dancoff equation:  $\sum_{p'h'} A_{ph,p'h'} X_{p'h'} = E X_{ph}$

$$A_{ph,p'h'} = (\epsilon_p - \epsilon_h) \delta_{ph,p'h'} + \langle ph' | \bar{v} | hp' \rangle$$


$$(\epsilon_{ph} - E) X_{ph} + \lambda D_{ph} \cdot T = 0 \quad T \equiv \sum_{ph} D_{ph}^* X_{ph}$$


$$X_{ph} = -\lambda \frac{D_{ph} T}{\epsilon_{ph} - E}$$


$$T = \sum_{ph} D_{ph}^* X_{ph} = -\lambda \sum_{ph} \frac{|D_{ph}|^2}{\epsilon_{ph} - E} T$$

or

$$\frac{1}{\lambda} = \sum_{ph} \frac{|D_{ph}|^2}{E - \epsilon_{ph}}$$


(TDA dispersion relation)


(separable interaction)


$$H\psi = E\psi; \quad \psi = \sum_i C_i \phi_i$$


$$\longrightarrow \sum_j H_{ij} C_j = E C_i; \quad H_{ij} = \langle \phi_i | H | \phi_j \rangle$$

suppose  $H_{ij} = \epsilon_i \delta_{i,j} + \lambda f_i^* f_j$  (separable form)


$$(\epsilon_i - E) C_i + \lambda f_i^* \underbrace{\sum_j f_j C_j}_{\equiv T} = 0$$


$$C_i = -\lambda \frac{T f_i^*}{\epsilon_i - E}$$


$$T = -\lambda \sum_j \frac{|f_j|^2}{\epsilon_j - E} T$$


$$\frac{1}{\lambda} = \sum_i \frac{|f_i|^2}{E - \epsilon_i}$$

# Graphical solutions

$$\frac{1}{\lambda} = \sum_{ph} \frac{|D_{ph}|^2}{E - \epsilon_{ph}}$$




Figure 8.4. Graphical solution of Eq. (8.18).

(note) in the degenerate limit:  $\epsilon_{ph} \sim \epsilon$

$$E = \epsilon_{ph} + \lambda \sum_{ph} |D_{ph}|^2, \quad |\nu\rangle = \sum_{ph} D_{ph} a_p^\dagger a_h |HF\rangle$$

*coherent superposition of 1p1h states*



Iso-scalar type modes:  $E < \epsilon_{ph} \rightarrow \lambda < 0$  (attractive)


Iso-vector type modes:  $E > \epsilon_{ph} \rightarrow \lambda > 0$  (repulsive)

### Experimental systematics:

IV GDR:  $E \sim 79 A^{-1/3}$  (MeV)  $\longleftrightarrow \epsilon_{ph} \sim 41 A^{-1/3}$

IS GQR:  $E \sim 65 A^{-1/3}$  (MeV)  $\longleftrightarrow \epsilon_{ph} \sim 82 A^{-1/3}$

(note) single particle potential:  $\hbar\omega \sim 41 A^{-1/3}$  (MeV)




$^{197}\text{Au}$

$E_{\text{GDR}} = 14$  (MeV)


$\epsilon_{ph} \sim 41 \cdot 197^{-1/3}$

$\sim 7$  (MeV)

$$\langle ph^{-1} | \bar{v} | p'h'^{-1} \rangle = \langle ph' | \bar{v} | hp' \rangle = \langle ph' | v | hp' \rangle - \langle ph' | v | p'h \rangle$$




=




Direct term

-



Exchange term

$$\left\{ \begin{array}{l} \langle PP^{-1} | \bar{v} | PP^{-1} \rangle \sim \langle NN^{-1} | \bar{v} | NN^{-1} \rangle = D - E \\ \langle PP^{-1} | \bar{v} | NN^{-1} \rangle = D \quad (\text{no charge exchange}) \end{array} \right.$$



$$\langle IS | \bar{v} | IS \rangle = 2D - E \sim D$$

$$\langle IV | \bar{v} | IV \rangle = -E \sim -D$$

$$|IS\rangle \propto |NN^{-1}\rangle + |PP^{-1}\rangle$$

$$|IV\rangle \propto |NN^{-1}\rangle - |PP^{-1}\rangle$$

# Random Phase Approximation

Tamm-Dancoff Approximation:  $|\nu\rangle = Q_\nu^\dagger |HF\rangle = \sum_{ph} X_{ph} a_p^\dagger a_h |HF\rangle$

(superposition of 1p1h states)

$$[H, Q_\nu^\dagger] \approx E_\nu Q_\nu^\dagger$$

$$\iff \langle HF | [\delta Q, [H, Q_\nu^\dagger]] | HF \rangle = E_\nu \langle HF | [\delta Q, Q_\nu^\dagger] | HF \rangle$$

Drawbacks:

➤ No influence of  $\nu$  in the ground state

$$E_{coll} = \epsilon + \lambda \sum_{ph} |D_{ph}|^2 \quad \longleftarrow \text{Interaction is essential in describing collective excitations}$$

➤ Energy Weighted Sum Rule is violated in TDA

➤ Admixture of the spurious modes with the physical excitation modes

HF  $\longleftrightarrow$  Broken Symmetries (CM localization, rotation,.....)


Restoration of broken symmetries  $\longrightarrow$  Goldstone mode  
(spurious motion)

A better approximation: **the random phase approximation (RPA)**

$$|\nu\rangle = Q_\nu^\dagger |0\rangle = \sum_{ph} \left( X_{ph} a_p^\dagger a_h - Y_{ph} a_h^\dagger a_p \right) |0\rangle$$

(superposition of 1p1h states)

smearing of Fermi surface due to the residual interaction (g.s. correlation)



# RPA on a schematic model

Separable interaction:  $\langle ph' | \bar{v} | hp' \rangle = \lambda D_{ph} D_{p'h'}^*$

$$\frac{1}{\lambda} = \sum_{ph} \frac{|D_{ph}|^2}{E - \epsilon_{ph}} - \frac{|D_{ph}|^2}{E + \epsilon_{ph}}$$

(RPA dispersion relation)

Cf. TDA dispersion relation:  $\frac{1}{\lambda} = \sum_{ph} \frac{|D_{ph}|^2}{E - \epsilon_{ph}}$




Figure 8.11. Graphical solution of the dispersion relation (8.135).

# Spurious motion in RPA

Mean-Field Approximation  $\longleftrightarrow$  Broken symmetries

- Center of mass localization (single center)
- Rotational motion

## Restoration of broken symmetries

$\longrightarrow$  Zero mode (Nambu-Goldstone mode)

**RPA**  $\langle HF | [\delta Q, [H, Q_\nu^\dagger]] | HF \rangle = E_\nu \langle HF | [\delta Q, Q_\nu^\dagger] | HF \rangle$

$\curvearrowright$  if  $[H, \hat{O}] = 0$

Then  $\hat{O}$  is a solution of RPA with  $E=0$

$$\hat{O} = \sum_{ph} (O_{ph} a_p^\dagger a_h + O_{hp} a_h^\dagger a_p)$$

$\curvearrowright$  The physical solutions are exactly separated out from the spurious modes.

# Comparison between Skyrme-(Q)RPA calculation and exp. data





photo-absorption  
cross section  
(GDR)

K. Yoshida  
and T. Nakatsukasa,  
PRC83('11)021304

## low-lying collective states




Low-lying excited states in even-even nuclei are collective excitations, and strongly reflect the pairing correlation and shell structure







# Single-particle transition (Weisskopf unit)

$$B(E\lambda : I_i \rightarrow I_{gs}) = \frac{(1.2A^{1/3})^{2\lambda}}{4\pi} \left( \frac{3}{\lambda + 3} \right)^2 (e^2 \text{fm}^{2\lambda})$$





M. Yamagami and Nguyen Van Giai, PRC69 ('04) 034301